

# Almonds



| Stage                                       | Dosage (l/ha)   | Application                                                                                                                              | Number of Applications | Outcome                                                                                                 |
|---------------------------------------------|-----------------|------------------------------------------------------------------------------------------------------------------------------------------|------------------------|---------------------------------------------------------------------------------------------------------|
| At planting                                 | 1 - 2% dilution | Drench (Bag/Open roots)                                                                                                                  | 1                      | Improve plant establishment and increase root development.                                              |
| Post planting                               | 1%              | Foliar spray during the spring and summer every 14 days<br>OR<br>2 - 5 l/ha through irrigation                                           | 4 - 6                  | Improve plant establishment and increase root development.                                              |
| Pink bud                                    | 4 - 6           | Foliar spray                                                                                                                             | 1                      | Increase nut set and crop retention.                                                                    |
| During bloom (30 - 50% bloom or full bloom) | 4 - 6           | Foliar spray                                                                                                                             | 1                      | Increase nut set and crop retention.                                                                    |
| Petal fall                                  | 4 - 6           | Foliar spray                                                                                                                             | 1                      | Increase nut set and crop retention.                                                                    |
| Post-harvest                                | 4 - 6           | Apply at post harvest time or the beginning of Autumn time, in 1 or 2 opportunities 14 days apart, by foliar spray or irrigation system. | 1 - 2                  | Support the second root flush growth to increase root tip numbers and improve nutrient uptake reserves. |

All recommendations are based on field experience and trial results worldwide. Genetics by environment (G X E) must be considered in all applications as this has an impact on plant growth and production. Please contact your local supplier/advisor for additional technical assistance. Each recommendation should be adapted to local conditions and/or optimised depending on crop varieties, soil, environment and fertilisation programs.